

AD-A154 238

CONSUMMATIVE EXPRESSIONS FOR SATELLITE DRAG
PERTURBATIONS IN A SPHERICAL (U) NAVAL SURFACE
WEAPONS CENTER DAHLGREN VA A D PARKS AUG 84
NSWC/TR-84-37 SBI-AD-F350 020

1/1

UNCLASSIFIED


F/G 22/1

NL

END

FILMED

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A154 230

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NSWC TR 84-37	2. GOVT ACCESSION NO. AD-A154230	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CONSUMMATIVE EXPRESSIONS FOR SATELLITE DRAG PERTURBATIONS IN A SPHERICALLY SYMMETRICAL ATMOSPHERE		5. TYPE OF REPORT & PERIOD COVERED FINAL
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) A. D. PARKS		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS NAVAL SURFACE WEAPONS CENTER (CODE K13) DAHLGREN, VA 22448		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS DEFENSE MAPPING AGENCY WASHINGTON, DC 20305		12. REPORT DATE AUGUST 1984
		13. NUMBER OF PAGES 21
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) satellite drag perturbations Lagrange planetary equations exponential atmosphere		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A compact formalism is developed that provides expressions for satellite aerodynamic drag perturbations when an exponentially varying atmospheric density distribution is assumed. This formalism is well suited for efficient manipulation and computation using the electronic computer.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

FOREWORD

Series expansion formulae have been applied to the Gaussian form of the Lagrange planetary equations in order to develop a compact formalism that represents satellite drag perturbations produced by an exponentially varying atmospheric density distribution. The compact result lends itself to facile manipulation, either manually by the user or automatically by the electronic computer. This document has been reviewed and approved by R. W. Hill.

Released by:

T. A. Clare

T. A. CLARE, Head
Strategic Systems Department

DTIC
ELECTE
S **D**
APR 25 1985
B



Accession For	
NTIS GMAI	<input checked="" type="checkbox"/>
NTIS DTD	<input type="checkbox"/>
NTIS DTD	<input type="checkbox"/>
Distribution/	
Availability Codes	
Dist	Avail. and/or Special
A-1	

CONTENTS

	<u>Page</u>
INTRODUCTION.	1
THE VARIATION OF THE ELEMENTS	2
INTEGRAND EXPANSION AND THE $J_{n,k}$ FUNCTION	7
THE CONSUMMATIVE DRAG PERTURBATION EQUATIONS.	10
REFERENCES.	13
DISTRIBUTION.	(1)

TABLE

<u>Table</u>	<u>Page</u>
1 EXPRESSIONS FOR FORMING THE \mathcal{C}_n MATRICES	12

INTRODUCTION

In recent years, much attention has been devoted to the analytic representation of artificial satellite aerodynamic disturbances produced by the earth's atmosphere¹⁻⁶. Such representations tend to generally address satellite orbits in the small eccentricity regime (e.g. $e \lesssim .2$), so that truncated expansions about the eccentricity can be used to simplify the complicated perturbation equations. Quite satisfactory results are produced in this way for a variety of assumed atmospheric density structures.

This report has been prepared to demonstrate that when one is concerned with an exponentially varying atmospheric density distribution, the aerodynamic perturbation equations can be written in a compact form without resorting to series truncation. This formalism lends itself to computational expediency, especially for applications using the electronic computer, and allows the user to readily inflate the expressions to any desired order in the orbital eccentricity.

The following sections of this report discuss in detail the derivation of this compact formalism. The Gaussian form of the planetary equations and their preliminary manipulation into a more utilizable configuration are presented in the next section. The operations used in the expansion process of the derivation are then discussed and, in the following section, are applied to the reconfigured planetary equations to yield the desired results.

THE VARIATION OF THE ELEMENTS

The Gaussian form of the equations describing the variation of the Keplerian element set a , e , i , ω , and Ω are well known and given by⁷

$$\dot{a} = 2a^2 \left(\frac{v}{h} \right) F_I \quad (1)$$

$$\dot{e} = \left(\frac{\sqrt{1-e^2} \sin E}{v} \right) F_R + \frac{2}{v} \left(\frac{\cos E}{1-e \cos E} \right) F_I \quad (2)$$

$$(\dot{i}) = \left(\frac{r \cos u}{h} \right) F_C \quad (3)$$

$$\dot{\Omega} = \frac{r \sin u}{h} \left(\frac{1}{\sin i} \right) F_C \quad (4)$$

and

$$\dot{\omega} = - \left(\frac{\cos E + e}{ev} \right) F_R + 2 \frac{\sqrt{1-e^2}}{ev} \left(\frac{\sin E}{1-e \cos E} \right) F_I - \dot{\Omega} \cos i \quad (5)$$

where u is the true argument of latitude defined as

$$u = \theta + \omega \quad (6)$$

θ is the true anomaly, and E is the eccentric anomaly. The quantities r , v , and h are the satellite radial distance, speed, and angular momentum, respectively, and are given by the following expressions:

$$r = a(1 - e \cos E) \quad (7)$$

$$v = \left(\frac{\mu}{a} \right)^{1/2} \left(\frac{1 + e \cos E}{1 - e \cos E} \right)^{1/2} \quad (8)$$

and

$$h = [\mu a(1 - e^2)]^{\frac{1}{2}} \quad (9)$$

where μ is the earth's gravitational constant.

The acceleration components F_i ($i = R, I, C$) appearing in the Gaussian equations are defined by the vector equation

$$\vec{F} = F_R \hat{R} + F_I \hat{I} + F_C \hat{C} \quad (10)$$

where

$$\hat{C} = \frac{\vec{r} \times \dot{\vec{r}}}{|\vec{r} \times \dot{\vec{r}}|} \quad (11)$$

$$\hat{I} = \frac{\dot{\vec{r}}}{|\dot{\vec{r}}|} \quad (12)$$

and

$$\hat{R} = \hat{I} \times \hat{C} \quad (13)$$

Here \vec{r} and $\dot{\vec{r}}$ are the satellite position and velocity vectors.

The drag force \vec{F}_D per unit satellite mass can be well represented by the equation

$$\vec{F}_D = - \frac{C_D A}{2m} \rho V_A \vec{V}_A \quad (14)$$

where C_D is the drag coefficient, A is the effective cross sectional area of the satellite normal to the direction of motion, m is the satellite mass, ρ is the local atmospheric density, and \vec{V}_A is the relative velocity given by

$$\vec{V}_A = \vec{V} - \vec{V}_a \quad (15)$$

this expression, \vec{v} is the satellite velocity and \vec{v}_a is the atmospheric rotational velocity. To a very good approximation the atmospheric drag acceleration components can be expressed as⁷

$$F_R = - \frac{C_D A}{2m} \alpha \left(\frac{a^5}{\mu} \right)^{\frac{1}{2}} e \rho V \omega_e \cos i \sin E \frac{(1 - e \cos E)^{\frac{3}{2}}}{(1 + e \cos E)^{\frac{1}{2}}} \dot{E} \quad (16)$$

$$F_I = - \frac{C_D A}{2m} \alpha a \rho V (1 + e \cos E)^{\frac{1}{2}} (1 - e \cos E)^{\frac{1}{2}} \dot{E} \quad (17)$$

$$F_C = - \frac{C_D A}{2m} \alpha \left(\frac{a^5}{\mu} \right)^{\frac{1}{2}} \rho V \omega_e \sin i (1 - e \cos E)^2 \cos u \dot{E} \quad (18)$$

where ω_e is the earth's angular rotation rate and

$$\alpha = 1 - \left[\frac{a^3 (1 - e^2)}{\mu} \right]^{\frac{1}{2}} \omega_e \cos i \quad (19)$$

Assume now that the atmospheric density varies exponentially with r and has the form

$$\rho = \rho_p \exp[\beta(r_p - r)]$$

$$\rho = \rho_p \exp[-\beta a e + \beta a e \cos E] \quad (20)$$

where ρ_p is the local atmospheric density at perigee and β is the inverse density scale height that is assumed to be constant. Using Equations (7), (8), (9), (16), (17), (18), and (20) along with the identities

$$\begin{aligned} \cos^2 u &\equiv \sin^2 \omega + \cos 2\omega \left(\frac{\cos E - e}{1 - e \cos E} \right)^2 \\ &\quad - (1 - e^2)^{\frac{1}{2}} \sin 2\omega \frac{\sin E (\cos E - e)}{(1 - e \cos E)^2} \end{aligned} \quad (21)$$

$$\sin u \cos u \equiv -\sin \omega \cos \omega + \sin 2\omega \left(\frac{\cos E - e}{1 - e \cos E} \right)^2 + (1 - e^2)^{\frac{1}{2}} \cos 2\omega \frac{\sin E (\cos E - e)}{(1 - e \cos E)^2} \quad (22)$$

in Equations (1) through (5), one finds that after much algebraic manipulation the Gaussian equations for the variation of the elements can be cast into the following forms:

$$\frac{da}{dE} = -a^2 \exp[\beta a e \cos E] (1 - e^2 \cos^2 E)^{-\frac{1}{2}} (1 + 2e \cos E + e^2 \cos^2 E) \quad (23)$$

$$\begin{aligned} \frac{de}{dE} = -\frac{\lambda}{2} \exp[\beta a e \cos E] (1 - e^2 \cos^2 E)^{-\frac{1}{2}} & \left\{ e(1 - e^2)^{\frac{1}{2}} \omega_e \cos i \left(\frac{a^5}{\mu} \right)^{\frac{1}{2}} \right. \\ & \left[1 - 2e \cos E - (1 - e^2) \cos^2 E + 2e \cos^3 E - e^2 \cos^4 E \right] \\ & \left. + 2a [\cos E + e \cos^2 E] \right\} \quad (24) \end{aligned}$$

$$\begin{aligned} \frac{di}{dE} = -\frac{\lambda}{2} \omega_e \sin i \left[\frac{a^5}{\mu(1 - e^2)} \right]^{\frac{1}{2}} \exp[\beta a e \cos E] (1 - e^2 \cos^2 E)^{-\frac{1}{2}} & \cdot \\ & \left\{ (\sin^2 \omega + e^2 \cos 2\omega) - 2e(\sin^2 \omega + \cos 2\omega) \cos E \right. \\ & + (1 - e^4) \cos 2\omega \cos^2 E + 2e^3(\cos 2\omega + \sin^2 \omega) \cos^3 E \\ & - e^2(\cos 2\omega + e^2 \sin^2 \omega) \cos^4 E + (1 - e^2)^{\frac{1}{2}} \sin 2\omega \sin E \cdot \\ & \left. (e - \cos E - e^3 \cos^2 E + e^2 \cos^3 E) \right\} \quad (25) \end{aligned}$$

$$\frac{d\Omega}{dE} = -\frac{\lambda}{2} \omega_e \left[\frac{a^5}{\mu(1-e^2)} \right]^{\frac{1}{2}} \exp[\beta a e \cos E] (1 - e^2 \cos^2 E)^{-\frac{1}{2}} \cdot$$

$$\left\{ \sin \omega \cos \omega [(2e^2 - 1) - 2e \cos E + 2(1 - e^4) \cos^2 E] \right.$$

$$+ 2e^3 \cos^3 E + e^2(e^2 - 2) \cos^4 E] - (1 - e^2)^{\frac{1}{2}} \cos 2\omega \sin E \cdot$$

$$\left. [e - \cos E - e^3 \cos^2 E + e^2 \cos^3 E] \right\} \quad (26)$$

$$\frac{d\omega}{dE} + \frac{d\Omega}{dE} \cos i = \frac{\lambda a}{2} \exp [\beta a e \cos E] (1 - e^2 \cos^2 E)^{-\frac{1}{2}} \sin E \cdot$$

$$\left\{ \left(\frac{a^3}{\mu} \right)^{\frac{1}{2}} \omega_e \cos i [e + (1 - 2e^2) \cos E - e(2 - e^2) \cos^2 E] \right.$$

$$\left. + e^2 \cos^3 E] - \frac{2(1 - e^2)^{\frac{1}{2}}}{e} (1 + e \cos E) \right\} \quad (27)$$

re

$$\lambda = \frac{C_D A}{m} \alpha \rho_o \exp [-\beta a e] \quad (28)$$

Finally, in order to obtain the changes produced in the orbital elements atmospheric drag during one orbital revolution, Equations (23) through (27) be integrated with respect to E over 2π radians; i.e.,

$$\begin{pmatrix} \Delta a \\ \Delta e \\ \Delta i \\ \Delta \Omega \\ \Delta \omega \end{pmatrix}_{\text{DRAG}} = \int_0^{2\pi} \frac{d}{dE} \begin{pmatrix} a \\ e \\ i \\ \Omega \\ \omega \end{pmatrix} dE \quad (29)$$

INTEGRAND EXPANSION AND THE $J_{n,k}$ FUNCTION

In this section, a novel method will be used to provide expansion formulae for Equation (29). As can be seen, the right-hand side of this equation is composed of sums of integrals of the form

$$R_n(\beta a e) = \int_0^{2\pi} \exp[\beta a e \cos E] (1 - e^2 \cos^2 E)^{-1/2} \cos^n E \, dE \quad (30)$$

The radical appearing in the integrand of R_n can readily be expanded using the Legendre polynomial generating function⁸

$$(1 - 2tx + t^2)^{-1/2} = \sum_{k=0}^{\infty} t^k P_k(x) \quad (31)$$

if one lets

$$x = 0 \quad (32)$$

and

$$t = ie \cos E \quad (i = \sqrt{-1}) \quad (33)$$

The $P_k(x)$ in Equation (31) is the Legendre polynomial.

When Equations (32) and (33) are used, then

$$\cos^n E (1 - e^2 \cos^2 E)^{-1/2} = \sum_{k=0}^{\infty} \binom{2k}{k} \left(\frac{e}{2}\right)^{2k} \cos^{2k+n} E \quad (34)$$

$$\binom{2k}{k} = \frac{(2k)!}{(k!)^2} \quad (35)$$

has been made of the equalities

$$p_\ell(0) = \begin{cases} \frac{(-1)^p}{2^\ell} \binom{2p}{p} & \text{for } \ell = 2p \\ 0 & \text{for } \ell = 2p + 1 \end{cases} \quad (36)$$

the relations⁹

$$\cos^{2k+n} E = \begin{cases} \left(\frac{1}{2^{2k+n}} \right) (2k+n)! \sum_{j=0}^{k+\frac{n}{2}} \frac{\epsilon_j \cos(2jE)}{(k+j+\frac{n}{2})! (k-j+\frac{n}{2})!} & \text{for } n \text{ even} \\ \left(\frac{2}{2^{2k+n}} \right) (2k+n)! \sum_{j=0}^{k+\frac{n-1}{2}} \frac{\cos[(2j+1)E]}{(k+j+\frac{n+1}{2})! (k-j+\frac{n-1}{2})!} & \text{for } n \text{ odd} \end{cases} \quad (37)$$

is Neumann's number defined as

$$\epsilon_j = \begin{cases} 1 & \text{for } j = 0 \\ 2 & \text{for } j \geq 1 \end{cases} \quad (38)$$

in Equation (34), one finds that Equation (30) may be written as

$$R_n(\beta ae) = \sum_{k=0}^{\infty} e^{2k} J_{n,k}(\beta ae) \quad (39)$$

In this expression,

$$J_{n,k}(\beta ae) = \frac{\pi(2k+n)!}{2^{4k+n-1}} \binom{2k}{k} \left\{ \Delta_{n,e} \sum_{j=0}^{k+\frac{n}{2}} \left[\frac{e_j}{(k+j+\frac{n}{2})!(k-j+\frac{n}{2})!} \right] \cdot \right. \\ \left. I_{2j}(\beta ae) + 2\Delta_{n,0} \sum_{j=0}^{k+\frac{n-1}{2}} \left[\frac{1}{(k+j+\frac{n+1}{2})!(k-j+\frac{n-1}{2})!} \right] \cdot \right. \\ \left. I_{2j+1}(\beta ae) \right\} \quad (40)$$

where

$$\Delta_{n,e} = \begin{cases} 1 & \text{for even } n \\ 0 & \text{for odd } n \end{cases} \quad (41)$$

$$\Delta_{n,0} = \begin{cases} 1 & \text{for odd } n \\ 0 & \text{for even } n \end{cases} \quad (42)$$

and $I_m(\beta ae)$ is a Bessel function of the first kind and imaginary argument defined by

$$I_m(\beta ae) = \frac{1}{2\pi} \int_0^{2\pi} \exp[\beta ae \cos E] \cos(mE) dE \quad (43)$$

THE CONSUMMATIVE DRAG PERTURBATION EQUATIONS

It should be noted that those portions of the right-hand sides of Equations (25), (26), and (27) that contain $\sin E$ multipliers, will vanish from the right-hand side of Equation (29), since

$$\int_0^{2\pi} \exp[\beta a e \cos E] \cos^m E (1 - e^2 \cos^2 E)^{-1/2} \sin E dE = 0 \quad (44)$$

Thus, Equations (30) and (39) may be used to express Equations (23) through (29) in the following compact matrix form

$$\underline{\Delta \varepsilon} = -\frac{\lambda}{2} \sum_{k=0}^{\infty} \sum_{n=0}^4 e^{2k+n} \underline{C}_n \underline{J}_{n,k} \quad (45)$$

where

$$\underline{\Delta \varepsilon} \equiv \begin{pmatrix} \Delta a \\ \Delta e \\ \Delta i \\ \Delta \Omega \\ \Delta \omega \end{pmatrix}_{\text{DRAG}} \quad (46)$$

$$\underline{C}_n \equiv \begin{pmatrix} \alpha_n^a & \beta_n^a & \gamma_n^a & \xi_n^a & \sigma_n^a \\ \alpha_n^e & \beta_n^e & \gamma_n^e & \xi_n^e & \sigma_n^e \\ \alpha_n^i & \beta_n^i & \gamma_n^i & \xi_n^i & \sigma_n^i \\ \alpha_n^\Omega & \beta_n^\Omega & \gamma_n^\Omega & \xi_n^\Omega & \sigma_n^\Omega \\ \alpha_n^\omega & \beta_n^\omega & \gamma_n^\omega & \xi_n^\omega & \sigma_n^\omega \end{pmatrix} \quad (47)$$

and

$$\mathcal{J}_{n,k} \equiv \begin{pmatrix} J_{n-2,k}(\beta a e) \\ J_{n-1,k}(\beta a e) \\ J_{n,k}(\beta a e) \\ J_{n+1,k}(\beta a e) \\ J_{n+2,k}(\beta a e) \end{pmatrix} \quad (48)$$

The elements of the matrix \mathcal{C}_n are generally functions of a , e , ω , and i and are presented in an easy to use format in Table 1.

TABLE 1. EXPRESSIONS FOR FORMING THE C_n MATRICES†

Element (ϵ) Coefficient	a	e	i	Ω	ω
α_n^ϵ	0	0	$\cos 2\omega(\delta_{n2} - \delta_{n4})Y$ **	$2(\delta_{n2} - \delta_{n4})Z$ ***	$-2(\delta_{n2} - \delta_{n4})Z \cdot \cos i$
β_n^ϵ	0	$X(\delta_{n1} - 2\delta_{n2} + \delta_{n3})$ *	0	0	0
γ_n^ϵ	$2a^2(\delta_{n0} + 2\delta_{n1} + \delta_{n2})$	0	$\sin^2 \omega(\delta_{n0} - 2\delta_{n1} + 2\delta_{n3} - \delta_{n4})Y - 2 \cos 2\omega(\delta_{n1} - \delta_{n3})Y$	$-\left[\delta_{n0} + 2(\delta_{n1} - \delta_{n3}) - \delta_{n4}\right]Z$	$\left[\delta_{n0} + 2(\delta_{n1} - \delta_{n3}) - \delta_{n4}\right] \cdot Z \cos i$
ξ_n^ϵ	0	$2a(\delta_{n0} + \delta_{n1}) + X(2\delta_{n2} - \delta_{n1} - \delta_{n3})$	0	0	0
σ_n^ϵ	0	0	$\cos 2\omega(\delta_{n0} - \delta_{n2})Y$	$2(\delta_{n0} - \delta_{n2})Z$	$-2(\delta_{n0} - \delta_{n2})Z \cdot \cos i$

†The δ_{ij} are Kronecker deltas
 $**Y = \omega_e \sin i \left[\frac{a^5}{\mu(1 - e^2)} \right]^{\frac{1}{2}}$

$*X = (1 - e^2)^{\frac{1}{2}} \omega_e \cos i \left(\frac{a^5}{\mu} \right)^{\frac{1}{2}}$
 $***Z = \omega_e \left[\frac{a^5}{\mu(1 - e^2)} \right]^{\frac{1}{2}} \sin \omega \cos \omega$

REFERENCES

1. Brouwer, D. and Hori, G., "Theoretical Evaluation of Atmospheric Drag Effects in the Motion of an Artificial Satellite," Astronomical Journal, Vol. 66, No. 5, June 1961, pp. 193-225.
2. King-Hele, D. G., Theory of Satellite Orbits in an Atmosphere, Butterworths Publishing Co., London, 1964.
3. Santora, F. A., "Satellite Drag Perturbations in an Oblate Diurnal Atmosphere," AIAA Journal, Vol. 13, No. 9, Sept. 1975, pp. 1212-1216.
4. Santora, F. A., "Drag Perturbations of Near-Circular Orbits in an Oblate Diurnal Atmosphere," AIAA Journal, Vol. 14, No. 9, Sept. 1976, pp. 1196-1200.
5. deLafontaine, J. and Garg, S. C., A Review of Satellite Lifetime and Orbit Decay Prediction, Institute for Aerospace Studies, University of Toronto, Toronto, Canada, June 1980.
6. Parks, A. D., Satellite Drag Perturbations in Nonsingular Variables Produced by an Oblate Diurnal Atmosphere, NSWC/DL TR (in preparation), Dahlgren, Virginia.
7. Fitzpatrick, P. M., Principles of Celestial Mechanics, Academic Press, Inc., New York, New York, 1970.
8. Messiah, A., Quantum Mechanics, Vol. 1, North-Holland Publishing Company, Amsterdam, Netherlands, 1965, pp. 492-494.
9. Oberhettinger, F., Fourier Expansions: A Collection of Formulas, Academic Press, New York, New York, 1973.

DISTRIBUTION

	Copies		Copies
Naval Space Command Dahlgren, VA 22448	5	Naval Research Laboratory ATTN: Mr. Al Bartholomew Washington, DC 20375	2
Library of Congress ATTN: Gift and Exchange Division Washington, DC 20540	4	Naval Oceanographic Office Bay St. Louis, MS 39522	2
Defense Mapping Agency ATTN: Mr. Jack Callander Washington, DC 20305	2	Office of Naval Research Physical Sciences Division 800 N. Quincy St. Arlington, VA 22217	2
Defense Mapping Agency Hydrographic/Topographic Center ATTN: Dr. Patrick Fell Washington, DC 20390	2	Air Force Geophysics Laboratory Hanscom Field Bedford, MA 01731	2
Defense Mapping Agency Aerospace Center ATTN: Dr. Robert Ballew St. Louis, MO 61118	2	Goddard Space Flight Center ATTN: Dr. David Smith Greenbelt, MD 20771	1
Office of Chief of Naval Operations Naval Oceanography Division (NOP-952) Bldg. 1, U. S. Naval Observatory Washington, DC 20390	2	Jet Propulsion Laboratory ATTN: Dr. William Melbourne Pasadena, CA 91103	1
Applied Research Laboratory University of Texas ATTN: Dr. Arnold Tucker Austin, TX 78712	1	The University of Texas at Austin ATTN: Dr. Byron Tapley Austin, TX 78712	1
Physical Sciences Laboratory New Mexico State University Box 3 - PSL ATTN: Dan Martin Las Cruces, NM 88003	1	Applied Physics Laboratory Johns Hopkins University Johns Hopkins Road ATTN: Harold Black Laurel, MD 20707	1
Office of Naval Operations Navy Space Systems Division (NOP-943) Washington, DC 20350	2		

DISTRIBUTION (Continued)

	Copies		Copies
Institute for Laboratory Astrophysics		Local:	
University of Colorado			
ATTN: Dr. Peter Bender	1	E31 (GIDEP)	1
Boulder, CO 80309		E411 (Hall)	1
		E431	10
U.S. Naval Observatory		F14	4
ATTN: W. L. Klepczaski	1	K05	1
D. D. McCarthy	1	K10	2
Washington, DC 20360		K12	5
		K13	35
Naval Space Surveillance System		K14	5
Dahlgren, VA 22448	2	K40	1
Headquarters Space Division (AFSC)			
Los Angeles Air Force Station			
Box 92960 Worldway Postal Center			
Los Angeles, CA 90009	2		

END

FILMED

6-85

DTIC